

Efficiency of ciprofloxacin removal by ozonation process with calcium peroxide from aqueous solutions

A. Rahmani*

J. Mehralipour**

A. Shabanlo**

S. Majidi**

*Professor of Environmental Health Engineering, Research Center for Health Sciences, Hamadan University of Medical Sciences, Hamadan, Iran

**M.Sc. Student of Environmental Health Engineering, School of Public Health, Hamadan University of Medical Sciences, Hamadan, Iran

*Abstract

Background: Advanced oxidation processes such as catalytic ozonation are efficient for the removal of antibiotics. Calcium peroxide is one of the catalysts that can be used as a source of hydrogen peroxide.

Objective: The aim of this study was to determine the efficiency of ciprofloxacin removal by ozonation process with calcium peroxide from aqueous solutions.

Methods: This experimental study was conducted in Hamadan University of Medical Sciences, 2013. The process consisted of ciprofloxacin with a concentration of 10-100 mg/L, pH=3-11 calcium peroxide with a concentration of 0-0.1 mg/L, 1 gr/min ozone (O₃), and 30 min reaction time in a semiconductor reactor.

Findings: In the optimal condition, at pH 3, calcium peroxide 0.025 mg/L, O₃ 1 gr/min, and initial antibiotic concentration of 10 mg/L, 92% of ciprofloxacin was removed and 45% of chemical oxygen demand was reduced.

Conclusion: With regards to the results, the ozonation process with calcium peroxide can be suitable for removal of ciprofloxacin contamination.

Keywords: Ciprofloxacin, Ozone, Calcium Peroxide

Citation: Rahmani A, Mehralipour J, Shabanlo A, Majidi S. Efficiency of ciprofloxacin removal by ozonation process with calcium peroxide from aqueous solutions. J Qazvin Univ Med Sci. 2015; 19 (2): 55-64.

Corresponding Address: Saba Majidi, Hamadan University of Medical Sciences, Shahid Fahmide Blvd., Hamedan, Iran

Email: saba.majidi44@yahoo.com

Tel: +98-919-3815592

Received: 1 JUN 2014

Accepted: 29 NOV 2014